

Entropy of Warped Taub-NUT AdS Black String via the Brick Wall Method

Chong Oh Lee

*Department of Physics, Kunsan National University,
Kunsan 573-701, Korea
cohlee@kunsan.ac.kr*

Abstract

When we consider five-dimensional warped Taub-NUT/bolt AdS black string with minimally coupled massive scalar field, we calculate entropies by using the brick wall method. It is found that they are proportional to being quadratically divergent in a cutoff parameter. In particular, we show that the entropy of warped Taub-NUT AdS black string holds for an area law in the bulk as well as on the brane.

1 Introduction

One can simply obtain p -brane solutions by adding a number of p extra dimensions to solutions for Einstein vacuum equations in d -dimensional spacetime [1, 2]. The simplest case of them can be given as black string solutions by adding just one extra dimension in higher dimensional spacetime. However, one cannot find asymptotically AdS black string solutions through an trivial way in as the above asymptotically flat case. They are able to be found after imposing warped geometry or working numerically [3, 4] obtained by solving Einstein equation with an extra dimension. It is of interest to find the black string solution in Taub-NUT AdS background.

The brick wall model was suggested in [5] search for black hole with a microscopic quantum states, and as the explanation of the origin of black hole entropy. It was investigated through the extensive application of various black holes [6]–[13].

In fact, black string forms when the matter field in five-dimensional spacetime is trapped on four-dimensional spacetime and undergoes gravitational collapse towards black hole. Comparing with horizon topology S^2 in four-dimensional black hole solution, its horizon topology is $S^2 \times S^1$ since such horizon extends into the extra dimension. Furthermore, the Bekenstein-Hawking entropy with the geometric character comes from the contributions of fields near the horizon. In this context, it was probed with black string [14]. In particular, it is shown that the entropy in Taub-NUT metric is quadratically divergent in a cutoff parameter [15] even if Taub-NUT metric has special property such as the solutions of the metric are not asymptotically flat (AF) but asymptotically locally flat (ALF) [16, 17]. Hence, it is of interest to investigate the corresponding situation in the case of warped Taub-NUT/bolt AdS black string.

On the other hand, it is well known that black string solution [18] is suggested in order to investigate how the physics of black holes is affected from the extra dimension with warped geometry in Randall-Sundrum (RS) brane world models [19, 20]. Such solution is nothing but RS brane world models with appearing as a Schwarzschild black hole on the brane. In this case, it is found that the entropy is proportional to the area of the event horizon on the brane and in the bulk [9, 10]. Therefore, an intriguing question is whether an area law is valid for warped Taub-NUT AdS black string.

The paper is organized as follows: In the next section, we will yield warped Taub-NUT AdS black string obtained by solving Einstein equation with a negative cosmological constant in five-dimensional spacetime. In section 3, we will calculate a free energy and an entropy through counting of the quantized modes obtained by using a WKB approximation and show how such entropy is expressed in terms of a cutoff parameter

and a black hole's area. Next we will explore the entropy in the case of Taub-bolt and show it is also proportional to being quadratically divergent in a cutoff parameter. In the last section, we will give our conclusion.

2 Warped Taub-NUT AdS Black String

Warped Taub-NUT black string in five-dimensional AdS spacetime is given as

$$ds_5 = a^2(z) \left[-f(r)(dt + 2n \cos(\theta)d\phi)^2 + \frac{dr^2}{f(r)} + (r^2 + n^2)(d\theta^2 + \sin^2(\theta)d\phi^2) \right] + l_5^2 dz^2, \quad (2.1)$$

which satisfies the Einstein equation with a negative cosmological constant $\Lambda_5 = -6/l_5$

$$G_{AB} - \Lambda_5 g_{AB} = 0. \quad (2.2)$$

Here, l_5 is the AdS radius, the warp factor $a(z) = \cosh(z)$, and

$$f(r) = \frac{r^2 - n^2 - 2Mr + l_5^{-2}(r^4 + 6n^2r^2 - 3n^4)}{r^2 + n^2}.$$

Taking $\cosh(z) = 1/\cos(\varphi)$, warped Taub-NUT black string in AdS_5 is rewritten as

$$ds_5 = \frac{l_5^2}{\cos^2(\varphi)} \left[\frac{1}{l_4^2} \left(-f(r)(dt + 2n \cos(\theta)d\phi)^2 + \frac{dr^2}{f(r)} + (r^2 + n^2)(d\theta^2 + \sin^2(\theta)d\phi^2) \right) + d\varphi^2 \right], \quad (2.3)$$

by introducing the relation between the cosmological parameter l_5 in the bulk and the cosmological parameter l_4 on the blane [1, 2]

$$l_5 = l_4 \cos(\varphi) \quad (2.4)$$

The metric (2.3) on locally constant $\varphi = \varphi_0$ slice is reduced to AdS Taub-NUT metric [21] localized on a brane

$$ds_4 = -f(r)(dt + 2n \cos(\theta)d\phi)^2 + \frac{dr^2}{f(r)} + (r^2 + n^2)(d\theta^2 + \sin^2(\theta)d\phi^2). \quad (2.5)$$

We consider a scalar field Φ with mass m propagating in five-dimensional spacetime under the background (2.1), which is described by five-dimensional Klein-Gordon equation as

$$\frac{1}{\sqrt{-\det(g_{\mu\nu})}} \partial_\mu \left(\sqrt{-\det(g_{\mu\nu})} g^{\mu\nu} \partial_\nu \Phi \right) - m^2 \Phi = 0 \quad (2.6)$$

Then we set the wave function

$$\Phi(t, r, \theta, \phi, z) = \Psi(t, r, \theta, \phi)\chi(z), \quad (2.7)$$

which leads to

$$\frac{1}{a^4(z)l_5^2}\partial_z\left(a^4(z)\partial_z\chi(z)\right) - (m^2 - a^2(z)\mu^2)\chi(z) = 0, \quad (2.8)$$

$$\begin{aligned} & \left(\frac{4n^2 \cos^2(\theta)}{(r^2 + n^2) \sin^2(\theta)} - \frac{1}{f(r)} \right) \partial_t \Phi - \frac{4n \cos(\theta)}{(r^2 + n^2) \sin^2(\theta)} \partial_t \partial_\phi \Phi \\ & + \frac{1}{r^2 + n^2} \partial_r \left((r^2 + n^2) f(r) \partial_r \Phi \right) + \frac{1}{(r^2 + n^2) \sin(\theta)} \partial_\theta \left(\sin(\theta) \partial_\theta \Phi \right) \\ & + \frac{1}{(r^2 + n^2) \sin^2(\theta)} \partial_\phi^2 \Phi - \mu^2 \Phi = 0, \end{aligned} \quad (2.9)$$

where μ can be interpreted as the effective mass on the brane by introducing picture borrowed from RS brane world models [19, 20]. Since there are no explicit time dependent terms in Eq.(2.9) one may take the stationary solutions as the following

$$\Psi(t, r, \theta, \phi) = e^{-i\omega t} e^{i\alpha\phi} R(r) Q(\theta), \quad (2.10)$$

and one from Eq.(2.9) can get

$$\frac{1}{\sin(\theta)} \partial_\theta \left(\sin(\theta) \partial_\theta Q(\theta) \right) + \left[l(l+1) - \frac{\alpha^2 + 2\alpha(2\omega n \cos(\theta)) + 4\omega^2 n^2}{\sin^2(\theta)} \right] Q(\theta) = 0, \quad (2.11)$$

$$\partial_r \left((r^2 + n^2) f(r) \partial_r R(r) \right) - \left[\left(\mu^2 - \frac{\omega^2}{f(r)} \right) (r^2 + n^2) + l(l+1) - 4\omega^2 n^2 \right] R(r) = 0, \quad (2.12)$$

where l is the degree of spherical harmonic. Using Wentzel-Kramers-Brillouin (WKB) approximation, the z -dependent wave number is given as

$$k_z^2 = l_5^2 (m^2 - a^2(z)\mu^2), \quad (2.13)$$

and the r -dependent wave number

$$k_r^2 = \frac{h^2(r, l, \omega)}{f^2(r)}, \quad (2.14)$$

with

$$h^2(r, l, \omega) = \left[\left(\frac{4n^2 f(r)}{r^2 + n^2} + 1 \right) \omega^2 - \left(\frac{l(l+1)}{r^2 + n^2} + \mu^2 \right) f(r) \right]. \quad (2.15)$$

3 Statistical Entropy

In this section, we will explicitly calculate the entropy through counting of the quantized modes for the z -dependent wave number (2.13) and the r -dependent wave number (2.14) in previous section.

The extra dimensional degeneracy factor n_z is obtained through the following the semi-classical quantization condition

$$\pi n_z = \int_0^{z_c} dz k_z(z, \mu), \quad (3.16)$$

and the radial degeneracy factor n_r

$$\pi n_r = \int_{r_h+\epsilon}^D dr k_r(r, l, \omega). \quad (3.17)$$

From Eqs. (2.13) and (3.16), we obtain

$$\frac{dn_z}{d\mu} = \frac{ml_5}{\pi\mu} \gamma(\mu), \quad (3.18)$$

with

$$\gamma(\mu) = \mathbf{E}_2\left(\sin^{-1}(a(z_c)), \frac{\mu^2}{m^2}\right) - \mathbf{F}_1\left(\sin^{-1}(a(z_c)), \frac{\mu^2}{m^2}\right) - \mathbf{E}_1\left(\frac{\mu^2}{m^2}\right) + \mathbf{K}_1\left(\frac{\mu^2}{m^2}\right), \quad (3.19)$$

where $\mathbf{E}_1$ is the complete elliptic integral, $\mathbf{E}_2$ the elliptic integral of the second kind, $\mathbf{F}_1$ the elliptic integral of the second kind, and $\mathbf{K}_1$ the complete elliptic integral of the first kind. Finally, in the WKB limit, the total number of quantum state with energy not exceeding ω is expressed as

$$g(\omega) = \int dg(\omega) = \int dl(2l+1) \int d\mu \left(\frac{dn_z}{d\mu} \right) \int \pi dn_r. \quad (3.20)$$

The free energy F of the scalar field at inverse temperature β is given as

$$\begin{aligned} \pi\beta F &= - \int_0^\infty d\omega \frac{\beta g(\omega)}{e^{\beta\omega} - 1} \\ &= -\beta \int_0^\infty \frac{d\omega}{e^{\beta\omega} - 1} \int dl(2l+1) \int \frac{ml_5}{\pi} \frac{\gamma(\mu)d\mu}{\mu} \\ &\quad \times \int_{r_h+\epsilon}^L \frac{dr}{f(r)} \sqrt{\left(\frac{4n^2 f(r)}{r^2 + n^2} + 1 \right) \omega^2 - \left(\frac{l(l+1)}{r^2 + n^2} + \mu^2 \right) f(r)}. \end{aligned} \quad (3.21)$$

Here, the reality condition of the free energy leads to the following limits for the remaining integrals:

$$0 \leq l \leq \frac{1}{2} \left[-1 + \sqrt{1 - 4(n^2 + r^2) \left\{ \left(\frac{4n^2 f(r)}{r^2 + n^2} + 1 \right) \frac{\omega^2}{f(r)} - \mu^2 \right\}} \right], \quad (3.22)$$

and

$$m \leq \mu \leq \sqrt{\frac{4n^2 f(r)}{r^2 + n^2}} + 1 \frac{\omega}{\sqrt{f(r)}}. \quad (3.23)$$

On the integrating over l , the free energy F is given as

$$F = -\frac{2ml_5}{3\pi} \int_0^\infty \frac{d\omega}{e^{\beta\omega} - 1} \int \frac{\gamma(\mu)d\mu}{\mu} \times \int_{r_h+\epsilon}^L dr \left(\frac{r^2 + n^2}{f^2(r)} \right) \left[\left(\frac{4n^2 f(r)}{r^2 + n^2} + 1 \right) \omega^2 - \mu^2 f(r) \right]^{\frac{3}{2}}. \quad (3.24)$$

The main contributions to the free energy, on carrying out the integrals over μ , are

$$F \approx -\rho(z_c) \frac{2ml_5}{3\pi} \int_0^\infty d\omega \frac{\omega^3}{e^{\beta\omega} - 1} \int_{r_h+\epsilon}^L dr \left(\frac{r^2 + n^2}{f^2(r)} \right) \quad (3.25)$$

with

$$\rho(z_c) = a(z_c) - 1, \quad (3.26)$$

which in the approximation of $L \gg r_h$ leads to

$$F \approx \left[-\frac{2\pi^3 m}{45\epsilon\beta^4} \frac{r_h^2(r_h^2 + n^2)}{(3(r_h^2 + n^2)l_5^{-2} + 1)^2} \right] \rho(z_c)l_5, \quad (3.27)$$

where the followings are used:

$$f(r) \approx (r - r_h) \left. \frac{df(r)}{dr} \right|_{r=r_h} = (r - r_h) (3(r_h^2 + n^2)l_5^{-2} + 1), \quad (3.28)$$

and

$$\int_0^\infty d\omega \frac{\omega^3}{e^{\beta\omega} - 1} = \frac{\pi^4}{15\beta^4}. \quad (3.29)$$

The entropy is given as

$$S = \beta^2 \left(\frac{\partial F}{\partial \beta} \right) = \left[\frac{8\pi^3 m}{45\epsilon\beta^3} \frac{r_h^2(r_h^2 + n^2)}{(3(r_h^2 + n^2)l_5^{-2} + 1)^2} \right] \rho(z_c)l_5. \quad (3.30)$$

The cutoff near the horizon is referred to as the brick wall. The physical distance between the brick wall and the horizon is

$$\int_{r_h}^{r_h+h} ds = \int_{r_h}^{r_h+\epsilon} \frac{dr}{\sqrt{f(r)}}, \quad (3.31)$$

which leads to

$$\epsilon = \left[\frac{3(r_h^2 + n^2)l_5^{-2} + 1}{4r_h} \right] h^2. \quad (3.32)$$

In terms of this covariant cutoff parameter h , the ultraviolet divergent part of the scalar field entropy is given by

$$S = \left[\frac{32\pi^3 m}{45h^2\beta^3} \frac{r_h^3(r_h^2 + n^2)}{(3(r_h^2 + n^2)l_5^{-2} + 1)^3} \right] \rho(z_c)l_5, \quad (3.33)$$

which is written in terms of the parameter M

$$S = \frac{4\pi^3 m}{45h^2\beta^3 M^2} \left[\frac{(r_h^2 - n^2 + l_5^{-2}(r_h^4 + 6n^2r_h^2 - 3n^4))^3}{(r_h^2 + n^2)^2((r_h^2 + n^2)^{-1} + 3l_5^{-2})^3} \right] \rho(z_c)l_5. \quad (3.34)$$

When taking $M = n + 4(2 + \sqrt{2})n^3l_5^{-2}$, r_h is evaluated as $(\sqrt{2} + 1)n$. Imposing the inverse temperature $\beta = 8\pi n$, the entropy in warped Tabu-NUT AdS spacetime is obtained as

$$S_{\text{AdS,NUT}} = \left[\frac{0.13n^2m}{(20.49n^2l_5^{-2} + 1)^3h^2} \right] \rho(z_c)l_5. \quad (3.35)$$

Taking $M = ((41\sqrt{41} + 365)n^2l_5^{-2} + 40)n/32$ in warped Tabu-bolt AdS spacetime, r_h is given by $(\sqrt{41} + 5)(n/4)$, and the entropy

$$S_{\text{AdS,bolt}} = \left[\frac{0.29n^2m}{(27.38n^2l_5^{-2} + 1)^3h^2} \right] \rho(z_c)l_5. \quad (3.36)$$

They reveal that the entropy in warped Tabu-Taub-NUT/bolt AdS spacetime has an explicit dependence on the inversely square of the cutoff parameter h .

Since the area of the horizon $\mathcal{A}_h$ on the brane located at $z = z_c$ is

$$\mathcal{A}_h = \int_0^{2\pi} \int_0^\pi \sqrt{g_{\theta\theta}g_{\phi\phi}} d\theta d\phi = 4\pi a^2(z_c)(r_h^2 + n^2), \quad (3.37)$$

the above entropy (3.35) is written as

$$S_{\text{AdS,NUT}} = \left[\left(\frac{\rho(z_c)}{16(\sqrt{2} + 1)\pi a^2(z_c)} \right) \left(\frac{0.13m}{(20.49n^2l_5^{-2} + 1)^3} \right) \left(\frac{l_5}{h^2} \right) \right] \mathcal{A}_h, \quad (3.38)$$

which leads to

$$S_{\text{AdS,NUT}} = \left[\left\{ \left(\frac{\rho(z_c)}{16(\sqrt{2} + 1)\pi a^2(z_c)} \right) \left(\frac{0.13ml_4 \cos(\varphi)}{(\frac{20.49n^2l_4^{-2}}{\cos^2(\varphi)} + 1)^3} \right) \right\} \left(\frac{1}{h^2} \right) \right] \mathcal{A}_h, \quad (3.39)$$

by using the relation (2.4). Evidently, the entropy on the brane located at $z = z_c$ satisfies the black hole area law since the curly-brackets quantity is much more smaller than the cutoff parameter h if we take h to be of the order of Planck length.

Introducing the invariant length of the black string along the z -direction

$$\int_0^{z_c} dz \sqrt{g_{zz}} = z_c l_5 \equiv \mathcal{R}, \quad (3.40)$$

from the entropy (3.38) we have in the limit of enough large $\mathcal{R}$

$$S_{\text{AdS,NUT}} \approx \left[\left(\frac{0.13 \rho(z_c) m}{16(\sqrt{2} + 1) \pi z_c a^2(z_c)} \right) \left(\frac{1}{h^2} \right) \right] \mathcal{A}_{\text{BS}}, \quad (3.41)$$

where the area of the black string horizon $\mathcal{A}_{\text{BS}}$ in the bulk is given as

$$\mathcal{A}_{\text{BS}} = \mathcal{A}_h \times \mathcal{R}. \quad (3.42)$$

It is also shown that the entropy of warped Taub-NUT AdS black string in the bulk holds for an area law from assumption of h with the order of Planck length.

One the other hand, employing thermal relation $\mathcal{E} = \partial_\beta(\beta F)$, the thermal energy is given as

$$\mathcal{E} = \left[\frac{2\pi^3 m}{15\epsilon\beta^4} \frac{r_h^2(r_h^2 + n^2)}{(3(r_h^2 + n^2)l_5^{-2} + 1)^2} \right] \rho(z_c) l_5. \quad (3.43)$$

Taking $r_h = (\sqrt{2} + 1)n$ and $\beta = 8\pi n$ and substituting in (3.32), we obtain the thermal energy in the case of Taub-NUT

$$\mathcal{E} = \left[\frac{(17\sqrt{2} + 24)mn}{3840(6(2 + \sqrt{2})n^2 l_5^{-2} + 1)^3} \right] \rho(z_c) l_5, \quad (3.44)$$

and the thermal energy in the case of Taub-bolt

$$\mathcal{E} = \left[\frac{(1057\sqrt{41} + 6765)mn}{960(3(5\sqrt{41} + 41)n^2 l_5^{-2} + 8)^3} \right] \rho(z_c) l_5, \quad (3.45)$$

by taking $r_h = (\sqrt{41} + 5)(n/4)$. It is shown that both thermal energies depend on the bulk parameter z_c along the z -direction and diverge much more slowly as n grows up.

4 Conclusion

Previous studies of quantized modes of scalar fields in Taub-NUT background [15] have indicated that the cutoff parameter h is determined by matching the statistical entropy with the gravitational entropy even if it is changed by the relative values of the parameters M and n .

Here we found the similar results for warped Taub-NUT AdS black string. Our evaluations of quantized modes of scalar fields in various warped black strings background were shown that the cutoff parameter has the Planck scale only by appropriate choosing the size of the AdS radius l_5 , in contrast to the results in as [19, 20]. We examined the correction to the statistical entropy due to the extra dimension with warped geometry and found the statistical entropy is still quadratically divergent in the cutoff parameter as in Taub-NUT metric [15]. Next, we showed that the entropy of warped Taub-NUT AdS black satisfies an area law in the bulk as well as on the brane. Finally, we investigated the thermal energy in warped Taub-NUT AdS/bolt black string and found that both thermal energies depend on the bulk parameter z_c along the z -direction and diverge much more slowly as n grows up.

Recently, an instability issue was covered for all topological spacetime through investigations of quasinormal modes of warped black strings with nontrivial topologies in AdS_5 [24]. In this context it would be of interest to study the stability of our warped black string solution.

It was shown that four-dimensional BTZ black string [25] still holds for the Gubser-Mitra conjecture [26], correlating the classical and thermodynamic instabilities of black strings. Intriguing issue would be to accomplish to test this conjecture for warped Taub-NUT AdS black string.

References

- [1] R. Gregory and R. Laflamme, Phys. Rev. Lett. **70**, 2837 (1993) [hep-th/9301052].
- [2] R. Gregory and R. Laflamme, Nucl. Phys. B **428**, 399 (1994) [hep-th/9404071].
- [3] K. Copsey and G. T. Horowitz, JHEP **0606**, 021 (2006) [hep-th/0602003].
- [4] R. B. Mann, E. Radu and C. Stelea, JHEP **0609**, 073 (2006) [hep-th/0604205].
- [5] G. 't Hooft, Nucl. Phys. B **256**, 727 (1985).

- [6] R. B. Mann, L. Tarasov and A. Zelnikov, *Class. Quant. Grav.* **9**, 1487 (1992).
- [7] A. Ghosh and P. Mitra, *Phys. Rev. Lett.* **73**, 2521 (1994) [hep-th/9406210, hep-th/9406210].
- [8] W. T. Kim, J. J. Oh and Y. -J. Park, *Phys. Lett. B* **512**, 131 (2001) [hep-th/0103147].
- [9] A. J. M. Medved, *Class. Quant. Grav.* **19**, 405 (2002) [hep-th/0110118].
- [10] H. K. Jassal and L. Sriramkumar, *Class. Quant. Grav.* **24**, 2589 (2007) [gr-qc/0611102].
- [11] S. Sarkar, S. Shankaranarayanan and L. Sriramkumar, *Phys. Rev. D* **78**, 024003 (2008) [arXiv:0710.2013 [gr-qc]].
- [12] B. S. Kay and L. Ortiz, arXiv:1111.6429 [hep-th].
- [13] K. S. Gupta, E. Harikumar, T. Juric, S. Meljanac and A. Samsarov, arXiv:1312.5100 [hep-th].
- [14] M. -l. Liu and H. -y. Liu, *Phys. Lett. B* **661**, 365 (2008) [arXiv:0811.0655 [gr-qc]].
- [15] K. Ghosh, *Phys. Rev. D* **67**, 124027 (2003) [gr-qc/0208055].
- [16] S. W. Hawking and C. J. Hunter, *Phys. Rev. D* **59**, 044025 (1999) [hep-th/9808085].
- [17] S. W. Hawking, C. J. Hunter and D. N. Page, *Phys. Rev. D* **59**, 044033 (1999) [hep-th/9809035].
- [18] A. Chamblin, S. W. Hawking and H. S. Reall, *Phys. Rev. D* **61**, 065007 (2000) [hep-th/9909205].
- [19] L. Randall and R. Sundrum, *Phys. Rev. Lett.* **83**, 3370 (1999) [hep-ph/9905221].
- [20] L. Randall and R. Sundrum, *Phys. Rev. Lett.* **83**, 4690 (1999) [hep-th/9906064].
- [21] A. Awad and A. Chamblin, *Class. Quant. Grav.* **19**, 2051 (2002) [hep-th/0012240].
- [22] R. B. Mann, *Phys. Rev. D* **60**, 104047 (1999) [hep-th/9903229].
- [23] R. Clarkson, L. Fatibene and R. B. Mann, *Nucl. Phys. B* **652**, 348 (2003) [hep-th/0210280].

- [24] S. Yin, B. Wang, R. B. Mann, C. O. Lee, C. -Y. Lin and R. -K. Su, Phys. Rev. D **82**, 064025 (2010) [arXiv:1004.4244 [hep-th]].
- [25] L. -h. Liu and B. Wang, Phys. Rev. D **78**, 064001 (2008) [arXiv:0803.0455 [hep-th]].
- [26] S. S. Gubser and I. Mitra, JHEP **0108**, 018 (2001) [hep-th/0011127].